

FRACTIONS

A DEFINING AND REPRESENTING FRACTIONS

Definition Rational Number

Any number that can be expressed as a **fraction** $\frac{a}{b}$, where a and b are integers and $b \neq 0$, is a **rational number**. The components of a fraction are formally defined:

$\frac{a}{b}$ ← **Numerator**: Indicates the number of parts being considered.
 ← **Denominator**: Indicates the total number of equal parts into which the unit has been divided.

A fraction can be represented in multiple ways.

- **Symbolic Form:**

$$\frac{2}{3}$$


- **Verbal Form:**

"Two-thirds" or "Two over three"

- **Linear Model:**



- **Number Line Model:**



B EQUIVALENT FRACTIONS

Definition Equivalent Fractions

Equivalent fractions are fractions that represent the same numerical value, even though they are written with different numerators and denominators.

The fundamental rule for generating equivalent fractions is to multiply or divide both the numerator and the denominator by the same non-zero number.


- When you multiply the numerator and the denominator by the same non-zero number (k), the fractions are equivalent:

$$\frac{a}{b} = \frac{k \times a}{k \times b}$$


- When you divide the numerator and the denominator by a common factor (k), the fractions are equivalent. This process is also known as **simplification**.

$$\frac{k \times a}{k \times b} = \frac{a}{b}$$

Ex:



Ex:



C SIMPLIFICATION

Definition Simplest Form of a Fraction

A fraction is considered to be in its **simplest form** (or lowest terms) when its numerator and denominator share no common factors other than 1.

Ex:

- The fraction $\frac{2}{3}$ is in simplest form because the only common factor of 2 and 3 is 1.
- The fraction $\frac{4}{6}$ is **not** in simplest form because 4 and 6 share a common factor of 2. It can be simplified to the equivalent fraction $\frac{2}{3}$.

Method Procedure for Simplifying a Fraction

To simplify a fraction, the numerator and denominator must be divided by their **Greatest Common Factor (GCF)**. This can be achieved by canceling all common factors.

Ex: Simplify the fraction $\frac{4}{6}$.

Answer:

$$\begin{aligned} \frac{4}{6} &= \frac{2 \times \cancel{2}}{3 \times \cancel{2}} \quad (\text{Cancel the common factor 2}) \\ &= \frac{2}{3} \end{aligned}$$

D CROSS MULTIPLICATION

Proposition Cross Multiplication Property

For any numbers a, b, c, d with $b \neq 0$ and $d \neq 0$,

$$\frac{a}{b} = \frac{c}{d} \quad \text{if and only if} \quad a \times d = b \times c$$

Ex: Solve x for $\frac{10}{5} = \frac{x}{8}$.

Answer:

$$\begin{aligned} \frac{10}{5} &= \frac{x}{8} \\ 5 \times x &= 10 \times 8 \quad (\text{cross multiplication}) \\ x &= 10 \times 8 \div 5 \quad (\text{dividing both sides by 5}) \\ x &= 16 \end{aligned}$$

E ADDITION AND SUBTRACTION

Definition Addition and Subtraction of Fractions with Common Denominators

For any numbers a, b, c with $c \neq 0$,

- To **add** fractions with common denominators, the numerators are added, and the denominator is retained.

$$\frac{a}{c} + \frac{b}{c} = \frac{a+b}{c}$$

- To **subtract** fractions with common denominators, the numerators are subtracted, and the denominator is retained.


$$\frac{a}{c} - \frac{b}{c} = \frac{a-b}{c}$$

Ex: Calculate $\frac{1}{4} + \frac{2}{4}$.

Answer:

•

$$\begin{aligned}\frac{1}{4} + \frac{2}{4} &= \frac{1+2}{4} \\ &= \frac{3}{4}\end{aligned}$$



Method Procedure for Adding or Subtracting Fractions

To add or subtract fractions with unlike denominators, follow this three-step procedure:

1. **Find a Common Denominator:** Identify a common multiple of the denominators.
2. **Create Equivalent Fractions:** Convert each fraction to an equivalent fraction with the common denominator.
3. **Add or Subtract the Numerators:** With the denominators now the same, perform the operation on the numerators and keep the common denominator.

Ex: Calculate $\frac{3}{4} + \frac{5}{6}$.


Answer:

- **Find a common denominator:** To add fractions, they must have the same denominator.

- Multiples of 4: 4, 8, **12**, 16, 20, ...
- Multiples of 6: 6, **12**, 18, 24, ...
- The smallest common denominator is **12**.

$$\begin{aligned}\frac{3}{4} + \frac{5}{6} &= \frac{3 \times 3}{4 \times 3} + \frac{5 \times 2}{6 \times 2} \\ &= \frac{9}{12} + \frac{10}{12} && \text{(common denominator = 12)} \\ &= \frac{9+10}{12} && \text{(adding numerators)} \\ &= \frac{19}{12}\end{aligned}$$

- **Visual representation:**



F MULTIPLYING A FRACTION BY A NUMBER

Definition Multiplication of a Fraction by an Number

To multiply a fraction by a number, multiply the number by the numerator of the fraction, and retain the original denominator.

For any numbers a, b, c with $c \neq 0$,


$$a \times \frac{b}{c} = \frac{a \times b}{c}$$

Ex: Calculate $3 \times \frac{2}{5}$.

Answer: Applying the definition, we multiply the integer (3) by the numerator (2) and keep the denominator (5).

$$3 \times \frac{2}{5} = \frac{3 \times 2}{5} = \frac{6}{5}$$

The calculation can be visualized as the repeated addition of $\frac{2}{5}$:



G MULTIPLICATION OF FRACTIONS

Definition Multiplication of Fractions

To multiply two fractions, one must multiply the numerators to find the numerator of the product, and multiply the denominators to find the denominator of the product.

For any numbers a, b, c, d with $b \neq 0$ and $d \neq 0$,

$$\frac{a}{b} \times \frac{c}{d} = \frac{a \times c}{b \times d}$$

Ex: Calculate $\frac{5}{2} \times \frac{3}{4}$.

Answer: Applying the rule for multiplication of fractions:

$$\frac{5}{2} \times \frac{3}{4} = \frac{5 \times 3}{2 \times 4} = \frac{15}{8}$$

Method Simplification by Canceling Common Factors

To simplify the multiplication process, any common factor that appears in both a numerator and a denominator can be canceled out before performing the multiplication. This is a direct application of simplification.

Ex: Calculate $\frac{31}{7} \times \frac{12}{31}$.

Answer: The number 31 is a factor in both a numerator and a denominator. It can be canceled before multiplying.

$$\begin{aligned}\frac{31}{7} \times \frac{12}{31} &= \frac{\cancel{31} \times 12}{7 \times \cancel{31}} \quad (\text{Cancel the common factor 31}) \\ &= \frac{12}{7}\end{aligned}$$

H DIVISION OF FRACTIONS

The operation of division is formally defined as the inverse of multiplication. To establish a procedure for dividing by a fraction, we must therefore utilize the concept of a multiplicative inverse. The multiplicative inverse is the value that "undoes" a multiplication, returning the result to the multiplicative identity, 1. This value is formally known as the **reciprocal**. Understanding the reciprocal is the necessary first step to developing the algorithm for fraction division.

Definition Reciprocal

For any non-zero number x , its **reciprocal** is the number which, when multiplied by x , yields the multiplicative identity, 1.

Proposition Reciprocal of a fraction

For a fraction $\frac{a}{b}$ (where $a \neq 0, b \neq 0$), its reciprocal is the fraction $\frac{b}{a}$.

Ex: State the reciprocal of $\frac{5}{7}$.

Answer: The reciprocal of $\frac{5}{7}$ is $\frac{7}{5}$.

Definition Division of fractions

To divide by a fraction, multiply by its reciprocal.
For any numbers a, b, c, d with $b \neq 0, c \neq 0$ and $d \neq 0$,

$$\frac{a}{b} \div \frac{c}{d} = \frac{a}{b} \times \frac{d}{c}$$

Equivalently, for a complex fraction:

$$\frac{\frac{a}{b}}{\frac{c}{d}} = \frac{a}{b} \times \frac{d}{c}$$

Ex: Calculate $\frac{2}{3} \div \frac{5}{7}$.

Answer: Applying the procedure for fraction division:

$$\begin{aligned}\frac{2}{3} \div \frac{5}{7} &= \frac{2}{3} \times \frac{7}{5} \quad (\text{Multiply by the reciprocal of the divisor}) \\ &= \frac{2 \times 7}{3 \times 5} \quad (\text{Multiply numerators and denominators}) \\ &= \frac{14}{15}\end{aligned}$$

I SIGN CONVENTIONS FOR FRACTIONS

Proposition Properties of Signs in Fractions

For any numbers a, b with $b \neq 0$,

1. A single negative sign results in a negative fraction. The sign can be placed with the numerator, the denominator,

or in front of the fraction.

$$\frac{-a}{b} = \frac{a}{-b} = -\frac{a}{b}$$

2. Two negative signs (one in the numerator and one in the denominator) result in a positive fraction, as the signs cancel each other out.

$$\frac{-a}{-b} = \frac{a}{b}$$

Ex: Simplify the fraction $\frac{-4}{-6}$.

Answer: The simplification follows these steps:

$$\begin{aligned}\frac{-4}{-6} &= \frac{4}{6} && \text{(Applying the sign rule: negative } \div \text{ negative} = \text{positive)} \\ &= \frac{2 \times \cancel{2}}{3 \times \cancel{2}} && \text{(Factoring and canceling the common factor of 2)} \\ &= \frac{2}{3}\end{aligned}$$

J FRACTIONS AS THE RESULT OF DIVISION

Proposition The Fraction as a Quotient

For any integers a and b (where $b \neq 0$), the division of a by b is represented by the fraction $\frac{a}{b}$.

$$a \div b = \frac{a}{b}$$

In this context:

- The **numerator** (a) corresponds to the **dividend**.
- The **denominator** (b) corresponds to the **divisor**.

Consequently, the fraction $\frac{a}{b}$ is the number which, when multiplied by the divisor b , yields the dividend a .

$$\frac{a}{b} \times b = a$$

Ex: The fraction $\frac{2}{3}$ is the same as saying "2 divided by 3".

$$2 \div 3 = \frac{2}{3}$$

The fraction $\frac{2}{3}$ is the number which, when multiplied by 3, gives 2:

$$\frac{2}{3} \times 3 = 2$$

K FRACTION AS A RATIO AND OPERATOR


Method From Ratio to Operation

To find a fraction of a quantity, multiply that quantity by the fraction:



$$\frac{a}{b} \text{ of } N = \frac{a}{b} \times N$$

Ex: In a class of 30 students, the ratio of girls to the total number of students is $\frac{2}{3}$ (i.e., two thirds of the class are girls). How many girls are there?

Answer: The fraction $\frac{2}{3}$ represents the part of the whole class that are girls.



- **Method 1 (unitary method).** First find one part: $30 \div 3 = 10$. Then take two parts: $10 \times 2 = 20$.



- **Method 2 (formula).**

$$\begin{aligned}
 \text{Number of girls} &= \frac{2}{3} \text{ of } 30 \\
 &= \frac{2}{3} \times 30 \\
 &= \frac{2 \times 30}{3} \\
 &= (2 \times 30) \div 3 \\
 &= 20.
 \end{aligned}$$

Check. $\frac{20}{30} = \frac{2}{3}$.

L FRACTIONS AS DECIMAL NUMBERS

Method Converting a Fraction to a Decimal Number

There are two primary methods for converting a fraction to its decimal equivalent.

- **Method 1: Direct Division**

Since a fraction $\frac{a}{b}$ is equivalent to the division $a \div b$, perform the division of the numerator by the denominator.

- **Method 2: Denominator as a Power of 10**

Find an equivalent fraction where the denominator is a power of 10 (e.g., 10, 100, 1000). The numerator of this new fraction can then be written as a decimal.

Ex: Convert $\frac{3}{4}$ to a decimal number.

Answer:

- **Applying Method 1 (Direct Division):**

$$\frac{3}{4} = 3 \div 4 = 0.75$$

$$\begin{array}{r}
 0.75 \\
 4 \overline{) 3.00} \\
 \underline{2.8} \\
 20 \\
 \underline{20} \\
 0
 \end{array}$$

- **Applying Method 2 (Power of 10):** We seek a number to multiply the denominator (4) by to get a power of 10. We know $4 \times 25 = 100$.

$$\frac{3}{4} = \frac{3 \times 25}{4 \times 25} = \frac{75}{100}$$

The fraction "seventy-five hundredths" is written as the decimal 0.75.

Method Converting a Decimal to a Fraction

The procedure for converting a terminating decimal to a fraction is as follows:

1. Write the decimal as the numerator of a fraction without the decimal point.
2. The denominator is 1 followed by as many zeros as there are decimal places in the original number.
3. Simplify the fraction to its lowest terms, if necessary.

Ex: Convert 1.3 to a fraction.

Answer:

- The number 1.3 has one decimal place.
- Write the number without the decimal point as the numerator: 13.
- The denominator will be 1 followed by one zero: 10.

The resulting fraction is $\frac{13}{10}$.

M REPRESENTING FRACTIONS GREATER THAN ONE

Definition Proper and Improper Fractions

Fractions are classified based on the relationship between the numerator and the denominator.

- A **proper fraction** is a fraction where the numerator is less than the denominator. Its value is always less than 1. Example: $\frac{2}{3}$.
- An **improper fraction** is a fraction where the numerator is greater than or equal to the denominator. Its value is always greater than or equal to 1. Example: $\frac{5}{3}$.

Definition Mixed Number

A **mixed number** is an alternative way to represent an improper fraction. It consists of an integer part (the number of whole units) and a proper fraction part.

The improper fraction $\frac{5}{2}$ can be visualized as two whole units and one half.

$$\frac{5}{2} = 2 + \frac{1}{2}$$

This is written as the mixed number $2\frac{1}{2}$. By convention, the addition sign is omitted:

$$2\frac{1}{2} \text{ is equivalent to } 2 + \frac{1}{2}$$

Caution: A mixed number like $2\frac{1}{2}$ *always* means $2 + \frac{1}{2}$, not $2 \times \frac{1}{2}$. If you mean multiplication, write $2 \times \frac{1}{2}$, $2 \cdot \frac{1}{2}$, or $(2)(\frac{1}{2})$.

N ORDER OF OPERATIONS

Proposition The Fraction Bar as a Grouping Symbol

In the order of operations (PEMDAS), the fraction bar functions as a grouping symbol for both the entire numerator and the entire denominator. This means that any expressions in the numerator and denominator must be fully evaluated to a single value before the final division, represented by the fraction itself, is considered.

The expression $\frac{A}{B}$ is mathematically equivalent to $(A) \div (B)$.

Therefore, the procedure is as follows:

1. Evaluate the entire expression in the numerator.
2. Evaluate the entire expression in the denominator.
3. Simplify the resulting fraction, if possible.

Ex: Simplify the expression $\frac{1+7}{3 \times 4}$.

Answer: Applying the order of operations for fractions:

$$\begin{aligned}\frac{1+7}{3 \times 4} &= \frac{(1+7)}{(3 \times 4)} && \text{(Recognize the implicit grouping)} \\ &= \frac{8}{12} && \text{(Evaluate the numerator and denominator)} \\ &= \frac{2 \times \cancel{4}}{3 \times \cancel{4}} && \text{(Simplify by canceling the common factor 4)} \\ &= \frac{2}{3}\end{aligned}$$