

# EQUATIONS OF LINES

There are many applications of vectors in geometry. While some of these applications can be addressed with other tools in 2-dimensional planar geometry, vector methods become particularly efficient and powerful in 3-dimensional space, especially when considering the relationships between lines and planes.


## A VECTOR EQUATION

The position of any point on a line can be described by a starting point and a direction of travel. In vector terms, this means that the position vector of any point on the line can be reached by starting with the position vector of a known point and adding a scalar multiple of the line's direction vector. This principle allows us to define a line in both two and three dimensions.

### Definition Vector Equation of a Line

The **vector equation of the line** is:

$$\vec{r} = \vec{a} + \lambda \vec{b}, \quad \lambda \in \mathbb{R}$$



## B PARAMETRIC EQUATIONS

Let a line pass through point  $A(a_1, a_2, a_3)$  with direction vector  $\vec{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$ . For any point  $R(x, y, z)$  on the line, the vector

equation  $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} a_1 \\ a_2 \\ a_3 \end{pmatrix} + \lambda \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$  leads to **parametric equations**:

$$\begin{cases} x = a_1 + \lambda b_1 \\ y = a_2 + \lambda b_2 \\ z = a_3 + \lambda b_3 \end{cases}, \quad \lambda \in \mathbb{R}$$

In 2 dimensions, the z-components are simply omitted.

### Definition Parametric Equations of a Line

The **parametric equations** of a line passing through a point  $A(a_1, a_2, a_3)$  with direction vector  $\vec{b} = \begin{pmatrix} b_1 \\ b_2 \\ b_3 \end{pmatrix}$  are given by the system:

$$\begin{cases} x = a_1 + \lambda b_1 \\ y = a_2 + \lambda b_2 \\ z = a_3 + \lambda b_3 \end{cases}$$

where the variable  $\lambda \in \mathbb{R}$  is the **parameter**.

## C CARTESIAN EQUATION IN PLANE

In two dimensions, a line can also be defined by a normal vector. Let  $\vec{n}$  be a normal vector to the line's direction, so that  $\vec{b} \cdot \vec{n} = 0$ . Starting from the vector equation  $\vec{r} = \vec{a} + \lambda \vec{b}$ , we take the scalar product of both sides with the normal vector  $\vec{n}$ :

$$\begin{aligned} \vec{r} \cdot \vec{n} &= (\lambda \vec{b} + \vec{a}) \cdot \vec{n} \\ \vec{r} \cdot \vec{n} &= \lambda \vec{b} \cdot \vec{n} + \vec{a} \cdot \vec{n} \quad (\text{distributivity}) \\ \vec{r} \cdot \vec{n} &= \vec{a} \cdot \vec{n} \quad (\vec{b} \cdot \vec{n} = 0) \end{aligned}$$

If we let  $\vec{r} = \begin{pmatrix} x \\ y \end{pmatrix}$  and the normal vector  $\vec{n} = \begin{pmatrix} a \\ b \end{pmatrix}$ , the expression  $\vec{r} \cdot \vec{n}$  becomes  $ax + by$ . The term  $\vec{a} \cdot \vec{n}$  is a constant, which we can call  $C$ . This leads to the familiar Cartesian form.

### Definition Cartesian Equation of a Line in 2D

The **Cartesian equation** of a line in 2D with normal vector  $\vec{n} = \begin{pmatrix} a \\ b \end{pmatrix}$  is given by:

$$ax + by = C$$

where  $C$  is a constant.

