


A OPTIMISATION

A.1 SOLVING OPTIMISATION PROBLEMS

Ex 1: A farmer fences off a rectangular field with a total of 4000 m of fencing. Since the field is located along a river, the farmer only needs to fence three of the four sides.




1. Let x be the length of each side perpendicular to the river, and y the side parallel to the river. Write an expression for the area A of the field in terms of x .
2. Determine the dimensions of the field which maximize the area (verification with the second derivative test is optional).


2. Find the value of x that maximizes the volume (verification with the second derivative test is optional).



Ex 3: Consider a semicircle of radius 20 cm. A rectangle is inscribed in the semicircle, with its base lying on the diameter.



Ex 2: A square sheet of paper 10 cm by 10 cm is made into an open box (no top) by cutting out squares of side x cm at each corner and folding up the sides.



1. Write an expression for the volume V of the box in terms of x .


1. Show that the area of the rectangle can be written as $A(x) = 2x\sqrt{400 - x^2}$, where x is half the base.
2. Find the value of x that maximizes $A(x)$, and determine the maximum area of the rectangle (verification with the second derivative test is optional).

B RATES OF CHANGE

B.0.1 SOLVING RELATED RATES PROBLEMS





Ex 5: Water is poured into an inverted right circular cone. The cone has a height of 12 cm and a radius of 6 cm at the top. The water is being poured in at a constant rate of 4 cm^3 per second. Let r be the radius of the water's surface and h be the height of the water at time t .



1. Show that the radius of the water's surface is always half of its height, i.e., $r = \frac{h}{2}$.
2. Find the rate at which the height of the water is increasing when the water is 8 cm deep.


Ex 4: A box has a volume of 1 m^3 , a square base, and an open top. Find the dimensions of the box which minimize its surface area (verification with the second derivative test is optional).





Ex 6: A ladder 10 m long is leaning against a vertical wall. The bottom of the ladder is pulled away from the wall at a rate of 0.5 m/s. Let x be the distance from the bottom of the ladder to the wall, and y the height of the top of the ladder on the wall.

1. Show that x and y are related by the equation $x^2 + y^2 = 100$.


2. Find the rate at which the top of the ladder is sliding down the wall when the bottom of the ladder is 6 m away from the wall.




Ex 8:  A street lamp is 5 m tall. A person 2 m tall walks away from the lamp at a speed of 1.2 m/s. Let x be the distance of the person from the base of the lamp, and s the length of their shadow.



1. Show that $\frac{5}{x+s} = \frac{2}{s}$.
2. Find the rate at which the length of the shadow is increasing when the person is 4 m away from the lamp.

Ex 7:  A spherical balloon is being inflated so that its volume increases at a constant rate of $100 \text{ cm}^3/\text{s}$. Let r be the radius of the balloon at time t .




1. Write the formula for the volume of the balloon in terms of r .
2. Find the rate at which the radius is increasing when $r = 5$ cm.



Ex 9: A gas is contained in a cylinder with a movable piston. The pressure P , volume V , and temperature T of the gas satisfy the ideal gas law:


$$PV = nRT,$$

where n and R are constants. The gas is heated so that the temperature increases at a constant rate of 2 K/s. At a certain instant, $T = 300$ K, $V = 0.01 \text{ m}^3$, and $P = 2.5 \times 10^5$ Pa. Assume the piston can move so that the pressure P remains constant.



1. Using the ideal gas law, express the volume V as a function of the temperature T (since P is constant).
2. Find the rate of change of the volume $\frac{dV}{dt}$ when $T = 300$ K.

Cooling



Ambient = 20°C

1. Use the data to determine the constant k .
2. Using Newton's law of cooling, find the value of $\frac{dT}{dt}$ when $T = 50^\circ\text{C}$.



Ex 10: A hot object is placed in a room where the ambient temperature is 20°C. Its temperature T decreases according to Newton's law of cooling:

$$\frac{dT}{dt} = -k(T - 20),$$

where k is a positive constant. At a certain moment, the object's temperature is 80°C and is cooling at a rate of 2°C per minute.

